

(43) International Publication Date 16 October 2003 (16.10.2003)

PCT

(10) International Publication Number WO 03/084323 A2

(51) International Patent Classification7:

A01N

- (21) International Application Number: PCT/US03/09441
- (22) International Filing Date: 27 March 2003 (27.03.2003)
- (25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

2002-096768

29 March 2002 (29.03.2002) Л

- (71) Applicant (for all designated States except US): SYN-GENTA PARTICIPATIONS AG [CH/CH]; Schwarzwaldallee 215, CH-4058 Basel (CH).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): HOMMA, Tamaki [JP/JP]; c/o Tottori University, Minami 4-101, Kozan-cho, Tottori, Tottori 680-8553 (JP). YODER, Joe [US/JP]; c/o Syngenta Crop Protection AG, Riehenstrasse 280, CH-4058 Basel (CH).

- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

 without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(I)

(54) Title: METHOD FOR INHIBITING DIFFERENTIATION AND FORMATION OF CONIFEROPHYTA MALE FLOWER BY TREATMENT WITH PROHEXADIONE COMPOUNDS

(57) Abstract: The present invention provides a method and composition for inhibiting the formation of Coniferophyta pollen, which has an excellent effect of inhibiting the formation of male Coniferophyta flowers without substantially inhibiting the elongation growth of the Coniferophyta trees. Prohexadione compounds are used as the active ingredients. The prohexadione compounds are, for example, as follows: formula (I).

0 03/084323 42